

An Observational Test of Dark Matter as Cold Fractal Clouds

Judith A. Irwin

irwin@astro.queensu.ca

Lawrence M. Widrow

widrow@astro.queensu.ca

and

Jayanne English[†]

english@astro.queensu.ca

Queen's University

Dept. of Physics

Kingston, Ontario, Canada, K7L 3N6

[†] Now at Space Telescope Science Institute

Received _____; accepted _____

ABSTRACT

Using the VLA, we have performed the first observational test of dark matter in the form of cold, primordial fractal clouds, as envisioned by Pfenniger, Combes, & Martinet (1994) and Pfenniger & Combes (1994). We show that, after a Hubble Time, primordial fractal clouds will convert most of their HI to H₂, but a small fraction of HI remains which is optically thick. This opens up a new window for detecting dark matter which may exist in this form. The detectability of such gas depends on its filling factor and temperature and therefore should be observable in absorption against a background source, with observations of sufficient sensitivity and resolution. The current VLA observations have made a first step towards this goal by taking advantage of a fortuitous alignment between the extension of the HI disk of the nearby galaxy, NGC 3079, and a background quasar, Q 0957+561. Our observations probe 28 independent beams against the quasar and all of velocity space between the extension of a flat rotation curve and a Keplerian decline for the halo region of NGC 3079. We do not detect any absorption features and investigate, in detail, the implication of this result for the hypothesis that dark matter is in the form of fractal clouds. In particular, we calculate the probability that our observations would have detected such clouds as a function of the model parameters. The chance of detection is significant for an interesting region of parameter space (fractal dimension $1.7 \lesssim D \lesssim 2$ and cloud radius $30 \text{ pc} < R_c < 3 \text{ kpc}$) and rises above 95% for a small region of parameter space. While our analysis does not rule out fractal clouds as dark matter, it does lay out the groundwork for future, more sensitive observations and we consider what form these might take to probe the range of possible cloud properties more deeply. It is interesting that the observations can rule out cold optically thin HI gas, if it exists, to a limit

of 0.001% of the dark matter. In contrast, the existence of cold HI in a fractal hierarchy would be an efficient way of hiding dark matter.

Subject headings: cosmology: dark matter — galaxies: halos, individual (NGC 3079) — quasars: absorption lines — ISM: clouds

1. Introduction

The question of extended disks around spiral galaxies, heavy halos, and indeed, the very nature of dark matter, has engaged astronomers for many years and dates to at least the early observations of flat rotation curves (e.g. Rubin, Ford, & Thonnard 1980). Since that time, observations too numerous to list have been carried out in attempts to search for evidence, direct or indirect, for this dark matter and to constrain its nature. Whatever the composition, there is now strong evidence that the dark halos surrounding spiral galaxies can extend to large distances. According to Ashman (1992), there is general agreement that the dark halo of the Milky Way, for example, extends to a radius between 100 kpc and 200 kpc.

It has been known for some time that primordial nucleosynthesis favors a density in baryons that is significantly greater than the density in luminous matter (e.g., Copi, Schramm, & Turner 1995). While recently, some of the results of primordial nucleosynthesis have been called into question, the conclusion that there are nonluminous baryons in the Universe remains firm and is in fact supported by other arguments from astrophysics and cosmology (Steigman, Hata, & Felten 1999). The question remains whether baryonic dark matter contributes significantly to halos in spiral galaxies. Baryonic dark matter has attracted considerable attention of late (see Carr 1994), especially in the light of recent reports of gravitational microlensing events toward the Large and Small Magellanic clouds, presumably (though not conclusively) due to massive compact halo objects (MACHOs) in the Milky Way (Alcock et al. 1997a, 1997b). At present, microlensing experiments are unable to fix the MACHO fraction in the Galaxy. In addition, the most probable mass for the lenses ($\sim 0.5 - 1M_{\odot}$) presents a puzzle since this rules out brown dwarfs and giant planets, initially considered to be the most plausible candidates for MACHOs.

A number of authors have considered cold gas as a dark matter candidate. Pfenniger, Combes, & Martinet (1994) and Pfenniger & Combes (1994) (hereafter PCM and PC) have

argued that the flat rotation curves of spiral galaxies might be accounted for entirely by cold, dense H_2 gas clouds. Henriksen & Widrow (1995) and Draine (1998) have suggested that very dense clouds might themselves act as gravitational lenses while Gerhard & Silk (1993, 1996) and de Paolis et al. (1995a, 1995b) have considered the possibility that gas might coexist with a population of MACHOs.

PCM and PC, in particular, have proposed that dark matter consists of cold primordial gas clouds in extended disks or halos around spiral galaxies where the gas within an individual cloud is distributed in a fractal structure. Far from sources of heating, the gas is thermalized to the 3 K background, assisted by small quantities of H_2 ice. PCM suggest that the slow accretion of gas toward the visible disk solves the “gas consumption problem” (i.e. that the timescale for gas consumption, as implied by current star formation rates, is shorter than the age of the galaxy) as well as the “disk-halo conspiracy” (i.e. the smooth, flat rotation curves through the transition region from disk to halo dominated rotation). Upon re-evaluation of the stellar recycling of gas, the former problem may not be as serious as previously thought (Kennicutt et al. 1994), but the latter problem remains. The PCM and PC proposal is motivated by fractal interpretations of the hierarchical structures observed over a wide range of scales for atomic and molecular interstellar clouds (see, e.g. Diamond et al. 1989, Vogelaar & Wakker 1994, and Elmegreen 1996). The HI in 7 members of the M 81 group (Westpfahl et al. 1999) and possibly in the Small Magellanic Cloud (SMC) (Stanimirovic et al. 1999) have also recently been shown to have a fractal structure. PCM suggest that the gas is in the form of 3 K molecular hydrogen which is unobservable directly and, if primordial, would also be unobservable via tracers such as CO. However, if a component of atomic hydrogen is present (see § 2.2), it could be detected in absorption against a background continuum source with observations of sufficient resolution and sensitivity.

In this paper we describe the first observational test of the PCM and PC model.

Specifically, we search for HI absorption associated with a foreground galaxy, NGC 3079, against the background quasar, Q 0957+561, which is fortuitously located nearby. Some preliminary results from this work have been presented in Irwin et al. (1999). In §2, we describe the PCM and PC model and discuss the neutral hydrogen content of the gas. In §3 we describe NGC 3079/Q 0057+561, the observations and data reduction, and in §4 we present the results. No significant absorption features are detected. The observations are extremely sensitive ($N(\text{HI}) = 1.4 \times 10^{17} \text{ cm}^2$) and indicate that no more than 0.001% of the dark matter halo can be in the form of cold *diffuse* HI. However, as we see in §5, we are unable to rule out fractal clouds as a dark matter candidate since the probability of detection is high ($> 95\%$) over a limited region of parameter space. This particular experiment lays the groundwork for future, more sensitive observations, as discussed in §6.

2. The Fractal Cloud Model

2.1. Elementary Cloudlets

The elementary building blocks in the model of PCM and PC are Jupiter-mass objects called “cloudlets” (“clumpuscles” by PC) which describe a natural minimum mass for fragmentation. This scale emerges by setting the free-fall time equal to the Kelvin-Helmholtz time and assuming virialization.

The cloudlet parameters, i.e. mass, radius, volume density, column density, and thermal line width, respectively, are given by:

$$M_* \simeq 4.0 \times 10^{-3} T^{1/4} \mu^{-9/4} f^{-1/2} M_\odot \quad (1)$$

$$R_* \simeq 1.5 \times 10^2 T^{-3/4} \mu^{-5/4} f^{-1/2} \text{ AU} \quad (2)$$

$$n_* \simeq 1.1 \times 10^8 T^{5/2} \mu^{3/2} f \text{ H cm}^{-3} \quad (3)$$

$$N_* \simeq 3.2 \times 10^{23} T^{7/4} \mu^{1/4} f^{1/2} \text{ H cm}^{-2} \quad (4)$$

$$v_* \simeq 9.1 \times 10^{-2} T^{1/2} \mu^{-1/2} \text{ km s}^{-1} \quad (5)$$

where T is the temperature, μ is the mean molecular weight, and f is a factor which accounts for departures from spherical symmetry and blackbody radiation (PC). Note that n_* is the mass density divided by the hydrogen mass, m_H (*not* by μm_H), i.e. $n_* \equiv 3M_*/4\pi R_*^3 m_H$, and therefore includes all forms of hydrogen as well as helium. The same holds for the surface density, N_* . These equations imply that warmer cloudlets are smaller, more massive, and thus more stellar-like.

For $T = 3 \text{ K}$, $0.1 \leq f \leq 1$ (see PC) and $1.3 \leq \mu \leq 2.3$ (primordial neutral gas), the resulting cloudlet parameters are, $M_* = 0.8\text{--}9.2 \times 10^{-3} M_\odot$, $R_* = 23\text{--}150 \text{ AU}$, $n_* = 0.25\text{--}6 \times 10^9 \text{ cm}^{-3}$, $N_* = 0.73\text{--}2.7 \times 10^{24} \text{ cm}^{-2}$, and $v_* = 0.10\text{--}0.14 \text{ km s}^{-1}$. Taking $\mu = 2.3$ (neutral $\text{H}_2 + \text{He}$, see § 2.2) and adopting $f = 1$, we have

$$M_* = 0.81 \times 10^{-3} M_\odot \quad (6)$$

$$R_* = 23 \text{ AU} \quad (7)$$

$$n_* = 6.0 \times 10^9 \text{ H cm}^{-3} \quad (8)$$

$$N_* = 2.7 \times 10^{24} \text{ H cm}^{-2} \quad (9)$$

$$v_* = 0.10 \text{ km s}^{-1} \quad (10)$$

2.2. Composition of the Cloudlets

Cloudlets at such high densities, if they exist, should consist only of neutral gas. While low density galactic HI disks are expected to be truncated by the extragalactic ionizing radiation field (Corbelli & Salpeter 1993, Maloney 1994, Dove & Shull 1994, see also Bland-Hawthorn et al. 1997), a simple application of the photoionization equilibrium

equation shows that the background UV photon rate at the current epoch ($4 \times 10^{-14} \text{ s}^{-1}$; Haardt & Madau 1996) is insufficient to produce any significant ionization in such high density cloudlets.

True primordial gas consists of atomic hydrogen since the “freeze-out” fraction of molecular hydrogen is only $[\text{H}_2/\text{H}] = 1.1 \times 10^{-6}$ (Galli & Palla 1998) and dust is not available to assist in the conversion from HI to H_2 . However, for these high density cloudlets, the 3-body reactions



may be important in converting HI to H_2 (see Palla et al. 1983). If we consider only these reactions, then the evolution of the number densities n_{HI} and n_{H_2} are governed by the equations:

$$\frac{dn_{\text{HI}}}{dt} = -2\mathcal{R}_1 n_{\text{HI}}^3 - 2\mathcal{R}_2 n_{\text{HI}}^2 n_{\text{H}_2} \quad (13)$$

$$\frac{dn_{\text{H}_2}}{dt} = \mathcal{R}_1 n_{\text{HI}}^3 + \mathcal{R}_2 n_{\text{HI}}^2 n_{\text{H}_2} \quad (14)$$

where $\mathcal{R}_1$ and $\mathcal{R}_2$ are the rate coefficients for reactions (11) and (12), respectively. These have not been measured at the temperatures and densities being considered here. Cohen and Westberg (1983) review the available data and recommend values, $\mathcal{R}_1 = 8.8 \times 10^{-33} \text{ cm}^6 \text{ s}^{-1}$, independent of temperature, T , and $\mathcal{R}_2 = 2.8 \times 10^{-31} (T/\text{K})^{-0.6} \text{ cm}^6 \text{ s}^{-1}$ above 80 K. In the absence of other data, we assume the above temperature dependencies hold down to 3 K so that for the case at hand, $\mathcal{R}_1$ is as given above and $\mathcal{R}_2 = 1.4 \times 10^{-31} \text{ cm}^6 \text{ s}^{-1}$.

When the cloudlets first form (time t_0), the gas is almost entirely in atomic form. However, on a timescale of order $(n_{\text{HI}}^2(t_0)\mathcal{R}_1)^{-1} \sim 10^5$ yrs, most of the hydrogen is converted to H_2 . Thereafter, the second reaction dominates and we have $n_{\text{H}_2} \simeq n_{\text{HI}}(t_0)/2$ and

$$n_{\text{HI}}(t) \sim \frac{1}{\mathcal{R}_2 n_{\text{HI}}(t_0) (t - t_0)} \quad (15)$$

For $n_{\text{HI}}(t_0) = 0.25 - 6 \times 10^9 \text{ cm}^{-3}$ (§ 2.1) and $(t - t_0) = 10$ Gyr, Eqn. (15) yields $n_{\text{HI}}(t) = 4.0 \times 10^3 - 9.5 \times 10^4 \text{ cm}^{-3}$. We take $n_{\text{HI}}(t)$ to represent the present density of HI in a cloudlet, i.e. $n(\text{HI})_*$. This implies that the fraction of the cloudlet in the form of HI, $f(\text{HI}) = n(\text{HI})_*/n_*$, is small, i.e. $f(\text{HI}) = 7 \times 10^{-7} - 2 \times 10^{-5}$.

If HI coexists with H_2 throughout the cloudlet, then the HI column density will be $N(\text{HI})_* = 2 n_{\text{HI}}(t) R_* = 2.8 \times 10^{18} - 4.3 \times 10^{20} \text{ cm}^{-2}$. Since in general,

$$\frac{N(\text{HI})}{[\text{cm}^{-2}]} = 1.82 \times 10^{18} \int \tau_v \frac{T_s}{[\text{K}]} \frac{dv}{[\text{kms}^{-1}]} \quad (16)$$

where τ_v is the HI optical depth at velocity, v , and T_s is the spin temperature (kinetic temperature) of the gas, the above parameters imply that $\tau \sim 5 - 700$, and hence even this small fraction of HI in the cloudlets would be optically thick. The detectability of such gas depends not on quantity, but on temperature and filling factor. Thus, even though HI may constitute a small fraction of the mass of the cloudlet, it can still provide a sensitive observational probe of the gas, if it exists.

2.3. Fractal Clouds

Cloudlets coalesce hierarchically in a fractal distribution to form larger structures called clouds (denoted with subscript, c , below). The clouds have parameters related to the parameters of the cloudlets via,

$$M_c = \mathcal{N}_c M_* = \left(\frac{R_c}{R_*} \right)^D M_* \quad (17)$$

$$n_c = \left(\frac{R_c}{R_*} \right)^{D-3} n_* \quad (18)$$

$$N_c = \left(\frac{R_c}{R_*} \right)^{D-2} N_* \quad (19)$$

$$v_c = \left(\frac{R_c}{R_*} \right)^{(D-1)/2} v_* \quad (20)$$

where $\mathcal{N}_c$ is the number of cloudlets in a cloud, v_c is the velocity dispersion of the cloud, R_c/R_* is the dynamic range of the fractal structure, and D is the *fractal dimension*. A pure diffuse smooth medium has $D = 3$ and lower values of D correspond to progressively lower filling factors. In general, we expect $D \lesssim 2$ since for $D > 2$, the clouds tend to dissolve due to internal collisions (PC). Molecular clouds in the ISM exhibit fractal structure with $1.4 \leq D \leq 2$. Recent results for HI in the M 81 group of galaxies give $1.2 \leq D \leq 1.5$ (Westpfahl et al. 1999) and for HI in the SMC, $D = 1.5$ (Stanimirovic et al. 1999).

3. Observations and Data Reduction

3.1. The NGC 3079 - Q 0957+561 Pair

A fortuitous alignment of the extension of the major axis of the galaxy, NGC 3079, with the background quasar, Q0957+561 has provided the opportunity to search for cold gas in an extended halo or an extended disk around the foreground galaxy (see Figure 1). NGC 3079 is at a distance of $D = 15.6$ Mpc ($H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$; $z = 0.004$) and has been mapped in HI by Irwin & Seaquist (1991) and Irwin et al. (1987). The outermost point at which emission is observed is denoted by a small star in Fig. 1. Q0957+561 is a gravitationally lensed system (Walsh et al. 1979) at a redshift of $z = 1.4$ and has also been extensively studied (e.g. Schmidt & Wambsganss 1998 and references therein). The

projected separation between the center of NGC 3079 (taken to be the kinematic center given by Irwin & Seaquist 1991), and Q0957+561 (taken to be the position of component A of RA (J2000) = $10^{\text{h}} 01^{\text{m}} 20^{\text{s}}.69$, DEC (J2000) = $55^{\circ} 53' 55''.9$) is 14.13 arcminutes, or 64.1 kpc at the distance of NGC 3079. For our purposes, we consider the lensed quasar to be simply a source of background radio continuum emission.

3.2. The Observations

Observations were carried out in the 21 cm spectral line on 03 January 1997 in the A configuration of the Very Large Array (VLA)¹. The flux calibrators were J1331+305 (3C286) and J0137+331 (3C48) which had flux densities at the central observing frequency of 14.79 Jy and 16.04 Jy, respectively, using the latest VLA calibration scale. The phase calibrator, J0957+553, for which we find a flux density of 2.45 ± 0.01 Jy, is separated from the source by only 43 arcminutes and was observed every ~ 20 minutes. All calibrators had UV restrictions which were applied during calibration. The field center was placed 13 arcseconds south of the quasar to avoid baseline-dependent map errors which tend to show up at the field center (Ekers 1989) and, for highest sensitivity, the 4IF mode was used.

The systemic velocity of NGC 3079 is 1117 km s^{-1} (heliocentric, optical definition) and the peak of the rotation curve is offset ± 215 from this value (Irwin & Seaquist 1991) with the background quasar occurring on the blueshifted side of the galaxy. Thus the central observing velocity was set to 974 km s^{-1} , which is between the velocity of a flat rotation curve for NGC 3079 extended to the projected distance of the quasar (i.e. 902 km s^{-1}) and the velocity expected at the position of the quasar for a Keplerian fall-off (1010 km s^{-1}). The bandwidth of 163.4 km s^{-1} encompasses both of these possibilities (though some

¹The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

channels at the band edges were lost due to a loss of sensitivity). At these frequencies, there should be no contamination by Galactic HI or by the lensing galaxy ($z = 0.36$) or known absorption systems ($z = 1.39$ and $z = 1.12$, Walsh et al. 1979) along the line of sight.

With on-line Hanning smoothing and 63 channels, the final channel width (= the spectral resolution) was 2.593 km s^{-1} . The primary flux density calibrators were also used as bandpass calibrators and were observed 4 times during the 10 hours of observations. The calibration observation closest in time to the source observation was applied. Examination of the average bandpass for each time showed that the maximum temporal change in the bandpass over the 4 scans was of order 0.2%.

UV data for each channel were Fourier Transformed and cleaned (e.g. Cornwell & Braun 1989) using uniform weighting (e.g. Sramek & Schwab 1989) (RA-DEC-Velocity) cube. A uniform weight clean continuum image was also made by averaging together the UV data for all channels (except the noisy ones at the band edge) before Fourier Transformation. The data were phase-only and then phase+amplitude self-calibrated (e.g. Cornwell & Fomalont 1989) until no improvement in the rms resulted. The final rms noise per channel is $0.61 \pm 0.01 \text{ mJy beam}^{-1}$, where the error represents the 1σ variation over all channels. This is ~ 1.3 times the theoretical rms for this weighting.

A continuum-subtracted cube was made by subtracting a linear fit to the visibilities over low-noise channels from each total emission channel. The resulting rms noise in the continuum-subtracted maps is $0.58 \pm 0.01 \text{ mJy beam}^{-1}$. Channel-to-channel variations due to the bandpass calibration are smaller than this. The spectral dynamic range, i.e. the ratio of peak continuum intensity to rms noise in the channel maps is $\sim 300/1$.

Naturally weighted cubes and continuum images were also made (with and without a UV taper), but since the rms was not significantly improved and the beam size was larger, subsequent analysis proceeded on the uniform weighting maps.

A summary of the observing and map parameters is given in Table 1.

4. Results

The continuum map is shown in Figure 2, with components labelled according to Greenfield et al. (1985) and Avruch et al. (1997). The lensing galaxy is associated with radio source, G, which is visible as a northern extension to component B. This map is very similar to the 18 cm image obtained by Avruch et al. (1997) from archival VLA data and confirms their discovery of sources R1, R2, and the northern source, N. Although not visible in Fig. 2, we also detect their southern source, S, in the naturally weighted map.

The continuum-subtracted cube was inspected for signs of emission or absorption features. No HI features were detected. In Fig. 3, we show spectra taken at the positions of the peaks of continuum components A, B, C, D, and E. The last panel shows the total flux as a function of velocity, integrated over a region in which the continuum emission is greater than $10 \times$ the continuum map rms noise value. These show no evidence for features above $\sim 3 \times$ the rms noise level per channel. Note that the noise in these spectra is dominated by the rms map noise, rather than by variations in the bandpass calibration which are smaller (§ 3.2). There is a slight hint of an absorption feature near zero velocity for components A, C, and E. However, this feature disappears if a spectrum is taken several pixels away from the continuum peaks but still within the same beam, and it also disappears if an average spectrum is taken over the beam at these positions. Moreover, if a spectrum is taken at a random position off-source, similarly sized features result. Thus, these features are only noise peaks.

We also formed 0th and 1st moment maps, i.e. total intensity and intensity-weighted mean velocity maps, respectively (not shown) in which low intensity emission was cut-off so that the higher intensity peaks would be highlighted. Again, there is no evidence for

emission or absorption in any channel.

In the event that HI exists at a low level over many channels, we also created a continuum-subtracted cube by doing linear fits to the visibilities over channels near the (low-noise) ends of the useable band, only. No evidence for any HI features could be seen in this cube either.

5. Constraints on Fractal HI in a Dark Matter Halo

5.1. Density of HI within NGC 3079

Our ability to detect clouds of the type proposed by PCM and PC depends both on the structure of individual clouds (the parameters R_c , R_* , and D) and the overall mass density in clouds. To quantify the latter, consider the hydrogen clouds in NGC 3079 to be smoothed into a continuous mass distribution throughout the halo of the galaxy. The line of sight to the background QSO intersects various galactocentric radii of this mass distribution. We define the hydrogen mass density at the radius closest to the center of NGC 3079 along this line of sight to be ρ_{system} .

Flat rotation curves observed in the outer parts of spiral galaxies suggest that dark matter halos follow an r^{-2} density law. Beyond this, little is known about the distribution of dark matter (e.g., shape of the halo). For definiteness, we assume an isothermal spheroid model for the halo of NGC 3079:

$$\rho_{\text{DM}} = \frac{\lambda(q)v_c^2}{4\pi G} \frac{1}{R^2 + z^2/q^2} \quad (21)$$

where v_c is the circular rotation speed, R is the galactocentric radius in the plane of the disk, z is the coordinate perpendicular to the plane ($r^2 = R^2 + z^2$), q is the flattening parameter ($q < 1$ for an oblate halo), and $\lambda(q) = \sqrt{1 - q^2}/(q \arccos q)$ is a geometric factor

equal to 1 for $q = 1$ and rising to 6.8 for $q = 0.1$. We have ignored the possibility of a core radius since we are only interested in the outer parts of the halo.

Roughly speaking, a maximum value $\rho_{\max}$ for ρ_{system} is obtained by assuming that the entire halo is composed of clouds (i.e., $\rho_{\text{system}} = \rho_{\text{DM}}$), that the halo is highly flattened ($q = 0.1$) and that the projected position of the quasar is in the equatorial plane of the halo ($R = 64 \text{ kpc}$; $z = 0$). Taking $v_c = 215 \text{ km s}^{-1}$ (Irwin & Seaquist 1991) we find $\rho_{\max} = 9.6 \times 10^{-26} \text{ g cm}^{-3}$. Let $F_H \equiv \rho_{\text{system}}/\rho_{\max}$ be the mean density in clouds relative to $\rho_{\max}$. $F_H = 1$ therefore corresponds to the “best-case” scenario for detecting clouds. Any possible effect which might reduce ρ_{system} is parametrized by F_H .

Integrating Eq. (21) from $-\infty$ to ∞ along the line of sight to the QSO and expressing the result in terms of a maximum surface density $N_{\max}$ (i.e., the surface density assuming $q = 0.1$ and $z = 0$) and the parameter, F_H , yields the following for the surface density:

$$N_{\text{system}} = N_{\max} F_H = \frac{\lambda(q) v_c^2}{4Gm_H R} F_H = 3.6 \times 10^{22} F_H \text{ H cm}^{-2} \quad (22)$$

The units, H cm^{-2} , are the same as in Eq. (4) and therefore represent the mass per unit surface area normalized by m_H , rather than the number of molecules per unit surface area. In deriving this expression, we have assumed that the density law is r^{-2} well beyond $R = 64 \text{ kpc}$. Obviously, one could introduce an additional parameter to allow for the possibility that the density law is steeper than r^{-2} in the outer parts of the halo.

5.2. Observational Parameters and Their Relation to Fractal Parameters

5.2.1. The Observational Parameters

For HI absorbing gas in front of a uniform background continuum source, the measured brightness temperature, T_B , in a given beam and velocity channel (i.e. a given resolution

element) is given by

$$T_B = [T_c e^{-\tau} + T_s(1 - e^{-\tau})] \mathcal{N} f_b f_v + T_c(1 - \mathcal{N} f_b f_v) \quad (23)$$

where T_c is the brightness temperature of the background continuum, T_s , τ , are the spin temperature and optical depth, respectively, of the gas, f_b , f_v are the filling factors for the beam area and velocity for a single cloud, and $\mathcal{N}$ is the number of clouds in the resolution element. Thus, the quantity, $\mathcal{N} f_b f_v$ is the filling factor for a resolution element. The factors, f_b and f_v , are dealt with separately since there are different conditions for saturating the beam and channel, respectively. The above equation reduces to the usual one for HI if this filling factor is unity. Aside from a constant of proportionality given by the Rayleigh-Jeans relation (T/I, Table 1), T_B is represented by the total emission channel maps.

After continuum subtraction,

$$\Delta T_B = (T_s - T_c)(1 - e^{-\tau}) \mathcal{N} f_b f_v \quad (24)$$

where ΔT_B represents the measured quantity in the continuum-subtracted cubes.

Since for most emission, $T_c \gg T_s$, then we can form the ratio,

$$\frac{\Delta T_B}{-T_c} = \begin{cases} \mathcal{N} f_v f_b & \text{for } \tau > 1 \text{ (fractal clouds)} \\ \tau \mathcal{N} f_v f_b & \text{for } \tau < 1 \text{ (optically thin gas)} \end{cases} \quad (25)$$

Figure 4 shows a map of the quantity $a \equiv \Delta T_B/(-T_c)$ for a single velocity channel where we have taken ΔT_B to be the negative of the rms map noise. To ensure that $T_c \gg T_s$ for all points, we have cut off all continuum emission less than $1.5 \text{ mJy beam}^{-1} = 583 \text{ K}$. Since the rms noise is constant over the map, this map has the appearance of the reciprocal of the continuum map. Such a map could be formed for any velocity channel, but the map

for any other channel would be identical, since both the continuum distribution and rms noise are the same, channel to channel. Note that this ratio map would be a map of upper limits to the optical depth under the more common assumptions of unity filling factor and optically thin gas.

Figure 4 provides observational constraints on the filling factor of optically thick fractal clouds through Eqn. (25). If a is the value read from Figure 4 for a particular resolution element, then our non-detection implies

$$\mathcal{N} f_b f_v < 3a \quad (26)$$

where we impose a 3σ upper limit, i.e. $[3a]$, on detectability.

The minimum of Figure 4 has a value 0.0033. Thus the strongest constraint we have for a single beam is $3a = 0.01$, i.e. less than 1% of this beam/velocity resolution element is filled with cold, fractal clouds. Of course, the clouds are distributed randomly and so we can tighten our constraint by considering the full data set. Our observations cover approximately 28 independent beams in any given channel and there are 163 channels² or 4564 independent beam/velocity resolution elements, in total.

5.2.2. Model Parameters

In this section, we apply the observational constraints to a family of fractal models defined by the parameters R_c/R_* , D , and F_H . We assume that the cloudlets are optically thick in HI (§2.2) and that their physical characteristics are given by Eqs.(6)-(10). We focus attention on values for D between 1 and 2 (§2.3) and values for R_c/R_* less than

²We have taken the full velocity coverage in this analysis, rather than the slightly reduced coverage due to the presence of a few noisy end channels.

3×10^7 which corresponds to a cloud radius $\simeq 3$ kpc. It is difficult to imagine a cloud 64 kpc from the center of the galaxy with a size much bigger than this. Indeed, such a cloud would subtend 2.7° if placed around the Milky Way, albeit with low covering factor.

For each point in parameter space, we calculate the probability that a cloud would have been detected. Recall that for a positive detection in a given resolution element, we require $\mathcal{N} f_b f_v > 3a$. It is straightforward to determine f_b and f_v as a function of R_c/R_* , D , and F_H (see below) while $\mathcal{N}$ can only be determined in a statistical sense. To see how this works, consider first a single beam. Let $P(\mathcal{N}_{\text{beam}})$ be the probability of finding $\mathcal{N}_{\text{beam}}$ clouds in this beam and $P(\mathcal{N}_{\text{beam}}|\mathcal{N}_{\text{eff}})$ be the conditional probability that at least one of its channels contain $\mathcal{N}_{\text{eff}}$ or more clouds, given $\mathcal{N}_{\text{beam}}$. By Bayes's theorem, the probability that at least one channel will contain $\mathcal{N}_{\text{eff}}$ or more clouds is

$$P(\mathcal{N}_{\text{eff}}) = P(\mathcal{N}_{\text{beam}})P(\mathcal{N}_{\text{beam}}|\mathcal{N}_{\text{eff}}) \quad (27)$$

The probability of having a detection in this beam is therefore

$$P = \sum_{\mathcal{N}_{\text{eff}}=\mathcal{N}_{\text{min}}}^{\infty} P(\mathcal{N}_{\text{eff}}) \quad (28)$$

where

$$\mathcal{N}_{\text{min}} = \frac{3a}{f_b f_v} \quad (29)$$

is the minimum number of clouds required for a detection.

It is straightforward to extend this analysis to all 28 beams. We use $j=1-28$ to label the different beams, i.e., P , a , and $\mathcal{N}_{\text{min}}$ all carry the subscript j . The probability for

detecting at least one cloud within the entire data set is

$$\mathcal{P} = 1 - \prod_j (1 - P_j) \quad (30)$$

The calculation of $\mathcal{P}$ requires, for each point in parameter space, f_v , f_b , $P(\mathcal{N}_{\text{beam}})$, and $P(\mathcal{N}_{\text{beam}}|\mathcal{N}_{\text{eff}})$. We determine these now, except for $P(\mathcal{N}_{\text{beam}}|\mathcal{N}_{\text{eff}})$ which is presented in the appendix.

Recall that the velocity dispersion of a cloud is v_c given by Eq. (20) while the width of a velocity channel is $\Delta v \equiv 2.59 \text{ kms}^{-1}$. A contour plot of v_c as a function of R_c/R_* and D is given in Figure 5. We estimate f_v by assuming that for $v_c \geq \Delta v$, the cloud covers the entire channel while for $v_c < \Delta v$ the cloud covers a fraction $v_c/\Delta v$ of the channel:

$$f_v = \min \{1, v_c/\Delta v\} \quad (31)$$

This ignores the obvious complication that a cloud can straddle more than one velocity channel. In particular, for $v_c \simeq \Delta v$, a cloud will generally cover some fraction of two neighboring channels.

The beam filling factor, f_b , depends on both the size of the cloud relative to the beam, and the area filling factor for the cloudlets in a cloud. If the cloud is larger than the beam ($R_c > R_b$ where $R_b = 47 \text{ pc}$ is the mean beam radius) then $f_b = (R_b/R_*)^{D-2}$, in analogy with Eq. (19). Conversely, if $R_c < R_b$ then $f_b = (A_c/A_b) (N_c/N_*) = (R_c/R_b)^2 (R_c/R_*)^{D-2}$ where A_c and A_b are the areas of the cloud and beam respectively. These results can be summarized in the following expression:

$$f_b = \left(\frac{R_{\min}}{R_b} \right)^2 \left(\frac{R_{\min}}{R_*} \right)^{D-2} \quad (32)$$

where $R_{\min} = \min(R_b, R_c)$. From Eqs. (29), (31) and (32) we construct a contour plot (Figure 6) for $\mathcal{N}_{\min}$ assuming $a = 0.0033$ (our most sensitive beam). From this figure, we can read off the number of clouds required in our most sensitive beam for a positive detection. For example, if the cloud system is described by the parameters $D = 1.64$ and

$R_c/R_* = 10^6$, then we would require at least one cloud in this beam for a positive detection. Likewise, for $D = 1.47$ and $R_c/R_* = 10^6$, we would require about 10 clouds in this one beam.

We now calculate $\langle \mathcal{N}_{\text{beam}} \rangle$, the expectation value for the number of clouds in a single beam. To do this, consider an area A large enough to contain a representative sample of clouds. The expected number of clouds in A will be $\mathcal{N}_A = (N_{\text{system}}/N_c)(A/A_c)$. Next consider a single beam in A . If $R_c > R_b$ there are three possibilities: cloud and beam do not intersect; cloud partially covers beam; cloud covers the entire beam. Likewise, the three possibilities for $R_c < R_b$ are: cloud and beam do not intersect; part of cloud covers beam; cloud entirely within beam. To avoid a rather awkward treatment that would attempt to take into account each of these possibilities, we use, as criteria for the cloud to be “in” the beam, that either the center of the beam is within the cloud perimeter (the $R_c > R_b$ case) or the center of the cloud is within the beam perimeter (the $R_c < R_b$ case). The probability for a single cloud in A to be also in the beam is therefore A_{max}/A where $A_{\text{max}} \equiv \max(A_b, A_c)$. The probability that none of the clouds in A overlap with the beam is $(1 - A_{\text{max}}/A)^{\mathcal{N}_A} \approx (1 - \mathcal{N}_A A_{\text{max}}/A)$. Since $A \geq \mathcal{N}_A A_{\text{max}}$, the expected number of clouds in the beam is then,

$$\langle \mathcal{N}_{\text{beam}} \rangle \simeq \mathcal{N}_A A_{\text{max}}/A = \frac{N_{\text{system}}}{N_c} \left(\frac{A_{\text{max}}}{A_c} \right) = \frac{F_H N_{\text{max}}}{N_*} \left(\frac{R_c}{R_*} \right)^{-D} \left(\frac{R_{\text{max}}}{R_*} \right)^2 \quad (33)$$

where $R_{\text{max}} \equiv \max(R_c, R_*)$.

A contour plot of $\langle \mathcal{N}_{\text{beam}} \rangle$ is given in Figure 7. We have assumed that $F_H = 1$, but the contours can be shifted proportionately for lower F_H according to Eqn (33). Note, however, that F_H approaching 1 is the only astrophysically interesting case.

The desired probability function, $P(\mathcal{N}_{\text{beam}})$ is given by the Poisson distribution:

$$P(\mathcal{N}_{\text{beam}}) = \frac{\langle \mathcal{N}_{\text{beam}} \rangle^{\mathcal{N}_{\text{beam}}} e^{-\langle \mathcal{N}_{\text{beam}} \rangle}}{\mathcal{N}_{\text{beam}}!} \quad (34)$$

The calculation of $P(\mathcal{N}_{\text{beam}}|\mathcal{N}_{\text{eff}})$ is more involved and therefore left to the appendix.

All of the ingredients are now in place and we can perform the calculation for $\mathcal{P}$, the probability for detecting at least one cloud in the entire dataset. Recall that an oblate halo filled with clouds, i.e. $F_H = 1$, is the best case scenario for detection. This is shown in Figure 8. The consequences of the beam area filling factor, Eq. (32), are evident in the plot. For example, clouds with $D < 1.7$ have $f_b \lesssim 0.02$ even if they fill the beam ($R_c > R_b$). Likewise, clouds with $R_c < 0.1R_b \simeq 10^{4.5}R_*$ have $f_b < 0.01$ even for $D = 2$. In these regions of parameter space one requires two or more clouds in a given resolution element which is unlikely. If fractal clouds exist in this range of parameter space, our observations would not have been able to detect them.

Our strongest constraints apply to the region, $1.75 \lesssim D \lesssim 1.85$ and $100 \text{ pc} \lesssim R_c \lesssim 3 \text{ kpc}$ ($R_c > R_b$) for which we find the highest probability, i.e. $\sim 95\%$. Thus, it is likely that if such fractal clouds exist, they would have been detected in our observations. A small cloud in this region, e.g. $D = 1.8$, $R_c = 110 \text{ pc}$, would have a mass, $5 \times 10^7 M_\odot$ (Eqn (17)) and a velocity dispersion of 25 km s^{-1} (Eqn (20)). There is another interesting region of parameter space which can be ruled out at the 90% probability level. This extends to much smaller clouds ($R_c < R_b$) with a higher covering factor (higher D). For example, $D = 1.9$, $R_c = 17 \text{ pc}$ (i.e. $\log(R_c/R_*)=5.2$) corresponds to a $6 \times 10^6 M_\odot$ cloud with a velocity dispersion of 22 km s^{-1} . Over all, there is a region of parameter space with a fractal dimension between about 1.7 and 2 and a fractal cloud dynamic range between about 10^5 and 10^7 within which there is a greater than 50% probability that a detection would have resulted from our observations.

5.2.3. Comparison with Optically Thin, Diffuse Gas

It is fruitful to examine the more conventional assumption that the gas is optically thin, diffuse, and that the velocity and beam filling factors are both 1. In this case, Fig. 4 provides upper limits to τ . Our strongest constraint, i.e. $\tau < 0.01$, then gives, for a single channel

(Eqn (16)), $N(HI) = 4.7 \times 10^{16} T_s \text{ cm}^{-2}$ and for $T_s = 3 K$, $N(HI) = 1.4 \times 10^{17} \text{ cm}^{-2}$, illustrating the high sensitivity of these absorption observations. This limit is deeper than previously measured 3σ noise values of $N(HI) = 1 \times 10^{18} (T_s/f_b) \text{ cm}^{-2}$ for Ly α absorption systems (Carilli et al. 1998), and is comparable to the upper limit of $3 \times 10^{16} T_s/f_b$ for HI absorption against more distant cooling flow galaxies (Dwarakanath et al. 1994). From Eqn (22), this implies that $F_H = 6.4 \times 10^{-6}$ for a flattened halo (and a factor of 7 higher for a spherical halo). Thus no more than 0.001% of the dark matter halo in NGC 3079 can be in the form of cold diffuse HI.

6. Discussion

The detection of dark matter remains one of the most ambitious quests in modern astrophysics. The observations described above demonstrate that one can search for a particular type of dark matter, cold, dense, hydrogen clouds, by looking for absorption features in the continuum spectrum of a background quasar. This is, of course, reminiscent of the rich study of Lyman- α clouds. However, in the case of NGC 3079/Q 0957+561, there is no doubt that the foreground absorber is a galaxy, in this case one which is nearby and has been well-studied in emission. We are also probing, in particular, the possible existence of halo gas far beyond the emitting HI disk which may be in an unusual form, i.e. cold and with a fractal structure.

The analysis indicates that for an all-cloud halo with model parameters in a limited though interesting range (fractal dimension $D > 1.7$; dynamic range of fractal $\sim 10^5 - 10^7$) there was a good chance for detection (see Fig. 8). The areas for improvement are obvious. Improved resolution would extend the region of sensitivity to lower values of R_c/R_* . Better signal to noise would bring probability levels up to where one would be able to rule out, at say the 99% confidence level, certain regions of parameter space. In addition, improved signal to noise would extend the region of sensitivity to lower values of the fractal dimension

D . Roughly speaking, we require $(R_b/R_*)^{D-2} > 3a$ where $a = (T_B - T_c)/(-T_c)$. An improvement in signal to noise by a factor of 10, at the same resolution, would allow us to probe clouds with D as low as 1.45.

Since these observations were at the highest resolution possible for connected arrays, the only practical improvements would be to search for objects like this in our own Galaxy, to use VLBI techniques, and/or to argue for more observing time for higher S/N. The difficulty in probing for objects in our own Galaxy is that the distance to the absorber may not be well known so it may not be possible to pin-point a particular absorber as a halo object. In moving to VLBI techniques, sensitivity is lost in comparison to connected arrays.

One possibility might be use VLBI techniques to look for individual Jupiter-mass cloudlets, directly, in our own Milky Way halo. Consider, for example, a small “sparse” cloud with $D = 1.3$ and $(R_c/R_*) = 10^3$ ($R_* = 23$ au, $R_c = 0.1$ pc), at a distance of 30 kpc which is distant enough to be far from Galactic sources of heating. An individual cloudlet will subtend an angle of 1.5 milliarcseconds while the cloud will subtend an angle, 1.5 arcseconds. The mean distance between clouds will be $(R_c^2/\mathcal{N}_c)^{1/2} = 1 \times 10^{-3}$ pc (see Eqn (17)), or 8 milliarcseconds. These parameters are favourable for multiple detections over a VLBI field of view against a strong background source. If the cloudlets are indeed optically thick at 3 K , the absorption signature should be obvious.

Indeed, VLBI HI absorption measurements have already detected a population of objects in the Galaxy with size scales of order 10s of au, as summarized and discussed by Heiles (1997). These tiny HI clouds, referred to by Heiles as “tiny scale atomic structure” (TSAS) have densities of $\sim 10^4$ cm $^{-3}$. This is in the range of $10^3 \rightarrow 10^5$ cm $^{-3}$ which we have estimated for the HI fraction of PC cloudlets (§2.2). TSAS features are relatively nearby as indicated by velocities, $|v| < 20$ km s $^{-1}$ (Diamond et al. 1989, Davis et al. 1996) or by absorption against pulsars which are within ~ 3 kpc (Frail et al. 1994). Consequently, they are near Galactic sources of heating (temperatures are often assumed to be ~ 50 K).

Given the sizes and densities of these objects, the possibility is raised that they are PC cloudlets which have already been accreted onto the Milky Way. A clear detection of CO associated with the HI absorbing systems would rule out this possibility.

To resolve the problem of unknown distance and argue that such objects are in the Galactic halo, a population of such objects would have to be detected at high velocities over a number of lines of sight.

7. Summary

Pfenniger et al. (1994) and Pfenniger & Combes (1994) have proposed a new model of dark matter consisting of cold clouds, radiatively coupled to the 3 *K* background, in an extended disk around galaxies. This gas is proposed to consist of virialized elementary cloudlets and to be of primordial composition except that at their high densities ($\sim 6 \times 10^9 \text{ cm}^{-3}$), gas phase reactions can convert most of the HI to H_2 . The cloudlets are arranged in a hierarchical fractal structure into larger clouds, the fractal structure being essential to ensuring that collisions do not dissipate the cloud and, in combination with low temperatures, that stars do not form.

In this work, we have shown that, at the temperatures and densities envisioned, a small fraction of HI will remain in the cloudlets and will likely be optically thick. Thus, their detectability depends on their temperature and the filling factor of the resolution element, rather than quantity. At a temperature of 3 *K*, the clouds cannot be seen in emission against the microwave background, but HI is potentially observable in absorption against a background source. This means that HI can be used as a probe of PC-type fractal clouds and opens a new window for exploring halo dark matter which may be in this form.

Using the VLA, we have carried out the first observational test of the PC model by taking advantage of a fortuitous alignment of a strong background quasar at a projected

separation of 64 kpc from the center of the nearby galaxy, NGC 3079. The quasar is aligned along an extension of the galaxy’s major axis. By tuning to the velocity of the foreground galaxy, we made 21 cm absorption measurements against the quasar, obtaining high spatial resolution ($1.2''$, or 28 beams across the background continuum) and high velocity resolution (2.6 km s^{-1}). We do not detect any HI absorption to a (minimum) 3σ upper limit $3 \Delta T_{B_v}/(-T_c) = 0.01$.

Our observational limits can be related to the filling factor of fractal clouds in a beam/velocity resolution element and we have considered, in detail, how the observations might constrain parameter space, defined by the quantities, R_c/R_* , D , and F_H , i.e. the dynamic range of the fractal structure, the fractal dimension, and the fraction of dark matter contained in such clouds, respectively. We find that much of parameter space cannot be ruled out by our observations. However, there is a region of parameter space (i.e., $1.7 \lesssim D \lesssim 2$ and $30 \text{ pc} \lesssim R_c \lesssim 3 \text{ kpc}$) within which there is a reasonable probability of detection. In the region $1.75 \lesssim D \lesssim 1.85$ and $100 \text{ pc} \lesssim R_c \lesssim 3 \text{ kpc}$, the probability of detection is as high as $\sim 95\%$ (see Fig. 8). Thus, it is unlikely that clouds with these parameters exist.

It is interesting that cold diffuse (optically thin) HI can be ruled out to a limit of 0.001% of the dark matter. In contrast, congregating the HI into optically thick fractal clouds is a very efficient way of hiding dark matter. Our analysis, applicable to any fractal clouds containing optically thick HI, is the first step in probing dark matter in this form. We outline possible ways of improving the observations in the future.

The authors wish to thank T. Abel, M. Mandy, R. Henriksen, N. Pelavas, and M. Walker for useful discussions. We have made use of the NASA/IPAC Extragalactic Database (NED). This work is supported by the Natural Sciences and Engineering Research Council of Canada

Appendix

In this appendix we calculate $P(\mathcal{N}_{\text{beam}}|\mathcal{N}_{\text{eff}})$, the probability that at least one channel in a given beam will contain $\mathcal{N}_{\text{eff}}$ or more clouds given $\mathcal{N}_{\text{beam}}$ clouds in this beam. The clouds are characterized by an internal velocity dispersion, v_c (assumed to be the same for all clouds) and a bulk velocity, $\bar{v}$ which is different for each of the clouds. For a given channel to contain contributions from two clouds, we require that their bulk velocities be separated by less than $\max(v_c - \Delta v, \Delta v)$. That is, for $v_c > 2\Delta v$, we require that the overlap of the clouds in velocity space should be greater than the width of a single channel whereas for $v_c < 2\Delta v$ we require that the separation of the clouds in velocity space be less than Δv . To continue, we need to know the distribution function for $\bar{v}$. In the absence of a detailed model for the cloud population, we assume that the $\bar{v}$ are equally likely to reside in any part of the 163 km s^{-1} velocity range covered by our observations. The probability that two particular clouds will contribute to a single channel is therefore $p = \max(v_c - \Delta v, \Delta v)/163 \text{ km s}^{-1}$. Put another way, we can think of there being $L \equiv 1/p$ bins in velocity space and we are asking for the probability that a certain number of clouds will appear in a single bin.

In general, if there are $\mathcal{N}_{\text{beam}}$ clouds in the beam, then the the probability that l_0 bins contain 0 clouds, l_1 bins contain one cloud, etc is given by

$$P(\mathcal{N}_{\text{beam}}, l_m, p) = p^{\mathcal{N}_{\text{beam}}} \frac{L!}{\prod_m l_m!} \frac{\mathcal{N}_{\text{beam}}!}{\prod_m (m!)^{l_m}} \quad (\text{A1})$$

The first factor is simply the probability of having a particular combination of $\mathcal{N}_{\text{beam}}$ clouds in L bins. The second factor handles the fact that we don't care which bins contain the different numbers of clouds. The final factor handles the fact that we don't care which clouds go into which bin. The desired probability is then given by

$$P(\mathcal{N}_{\text{beam}}|\mathcal{N}_{\text{eff}}) = \sum P(\mathcal{N}_{\text{beam}}, l_m, p) \quad (\text{A2})$$

where the sum is over all combinations of l_m consistent with the constraints that $\sum_m l_m = 1/p$, $\sum_m l_m m = \mathcal{N}_{\text{beam}}$ and $l_M \geq 1$.

To illustrate how the calculation will go, we consider a few simple cases. Clearly $P(\mathcal{N}_{\text{beam}}|1) = 1$ since the probability of having at least one channel with one or more clouds is unity provided $\mathcal{N}_{\text{beam}} \geq 1$. As discussed above, and in agreement with Eq A1 and A2, $P(2|2) = p$. For $P(4|2)$, there will be four contributions to Eq. A2 corresponding to $(l_0, l_1, l_2, l_3, l_4) = (1, 2, 1, 0, 0), (2, 1, 0, 1, 0), (3, 0, 0, 0, 1), (2, 0, 2, 0, 0)$. For the special case where $l = 1/p = 4$ we find $P(4|2) = 29/32$. This case is easy to check since the probability of having two or more clouds in one of the four channels is simply $1 - (\text{probability of having one cloud in each channel}) = 1 - 4!/4^4 = 29/32$.

REFERENCES

- Alcock et al. 1997a, ApJ, 486, 697
- Alcock et al. 1997b, ApJ, 497, L11
- Ashman, K. M. 1992, PASP, 104, 1109
- Avruch, I. M., Cohen, A. S., Lehár, J., Conner, S. R., Haarsma, D. B., & Burke, B. F. 1997, ApJ(letters), 488, L121
- Bland-Hawthorn, J., Freeman, K., & Quinn, P. J. 1997, ApJ, 490, 143
- Carr, B. 1994, ARA&A32, 531
- Carilli, C. L., Menten, K. M., Reid, M. J., Rupen, M. P., & Yung, Min Su 1998, ApJ, 494, 175
- Cohen, N., and Westberg, K. R. 1983, J. Phys. Chem. Ref. Data, 12, 560
- Copi, C. J., Schramm, D. N., & Turner, M. S. 1995, Science, 267, 192
- Corbelli, E., & Salpeter, E. E. 1993, ApJ, 419, 94
- Cornwell, T., & Braun, R. 1989 in ASP Conf. Series 6, Synthesis Imaging in Radio Astronomy, ed. R. A. Perley, F. R. Schwab, & A. H. Bridle, p. 167
- Cornwell, T., & Fomalont, E. B. 1989 in ASP Conf. Series 6, Synthesis Imaging in Radio Astronomy, ed. R. A. Perley, F. R. Schwab, & A. H. Bridle, p. 185
- Davis, R. J., Diamond, P. J., & Goss, W. M. 1996, MNRAS, 283, 1105
- Deguchi, S., & Watson, W. D. 1985, ApJ, 290, 578
- Diamond, P. J., Goss, W. M., Romney, J. D., Booth, R. S., Kalberla, P. M. W., & Mebold, U. 1989, ApJ, 347, 302

- Dove, J. B., & Shull, J. M. 1994, *ApJ*, 423, 196
- Dwarakanath, K. S., van Gorkom, J. H., & Owen, F. N. 1994, *ApJ*, 432, 469
- Ekers, R. D. 1989, in *ASP Conf. Series 6, Synthesis Imaging in Radio Astronomy*, ed. R. A. Perley, F. R. Schwab, & A. H. Bridle, p. 199
- Imegreen, B. 1996, *ApJ*, 471, 816
- Filippenko, A. V., & Sargent, W. L. W. 1992, *AJ*, 103, 28
- Frail, D. A., Weisberg, J. M., Cordes, J. M., & Mathers, C. 1994 *ApJ*, 436, 144
- Galli, D., & Palla, F. 1998, *Å*, 335, 403
- Greenfield, P. E., Roberts, D. H., & Burke, B. F. 1985, *ApJ*, 293, 370
- Haardt, F., & Madau, P. 1996, *ApJ*, 461, 20
- Heiles, C. 1997, *ApJ*, 481, 193
- Henriksen, R. N., & Widrow, L. W. 1995, *ApJ*, 441, 70
- Irwin, J. A. 1995, *PASP*107, 715
- Irwin, J. A., & Seaquist, E. R. 1988, *ApJ*, 335, 658
- Irwin, J. A., & Seaquist, E. R. 1990, *ApJ*, 353, 469
- Irwin, J. A., & Seaquist, E. R. 1991, *ApJ*, 371, 111
- Irwin, J. A., Seaquist, E. R., Taylor, A. R., & Duric, N. 1987, *ApJ*, 313, L91
- Irwin, J. A., Widrow, L. M., & English, J. 1999, *PASA*, in press
- Kennicutt, R. C. Jr., Tamblyn, P., & Congdon, C. W. 1994, *ApJ*, 435, 22
- Maloney, P. 1993, *ApJ*414, 41

- Palla, F., Salpeter, E. E., & Stahler, S. W. 1983, *ApJ*, 271, 632
- Pfenniger, D., & Combes, F. 1994, *A&A*, 285, 94 (PC)
- Pfenniger, D., Combes, F., & Martinet, L. 1994, *A&A*, 285, 79 (PCM)
- Rubin, V. C., Ford, W. K., & Thonnard, N. 1980, *ApJ*, 238, 471
- van Gorkom, J. H. 1993, in *The Environment and Evolution of Galaxies*, ed. J. M. Shull and H. A. Thronson, Jr. (Dordrecht, Kluwer), p. 345
- Schmidt, R., & Wambsganss, J. 1998, *A&A*, 335, 379
- Steigman, G., Hata, N., & Felten, J. E. 1999, *ApJ*, 510, 564
- Sramek, R. A., & Schwab, F. R. 1989 in *ASP Conf. Series 6, Synthesis Imaging in Radio Astronomy*, ed. R. A. Perley, F. R. Schwab, & A. H. Bridle, p. 185
- Stanimirovic, S., Staveley-Smith, L., Dickey, J. M., Sault, R. J., & Snowden, S. L. 1999, *MNRAS*, 302, 417
- Vogelaar, M. G. R., & Wakker, B. P. 1994, *Å*, 291, 557
- Walsh, D., Carswell, R. F., & Weymann, R. J. 1979, *Nature*, 279, 381
- Westpfahl, D. J., Coleman, P. H., Alexander, J., & Tongue, T. 1999, *AJ*, 117, 868

Fig. 1.— Digitized Sky Survey optical image of the foreground galaxy, NGC 3079, and the background quasar, Q 0957+561 (large star). The small star marks the approximate boundary of the HI emission associated with NGC 3079 as measured by Irwin et al. (1987).

Fig. 2.— Uniformly weighted continuum map of Q0957+561. Contours are at -0.26, 0.26 (2σ), 0.3, 0.4, 0.5, 0.8, 1.5, 2.5, 5, 10, 20, 30, 50, 75, 120, and 170 mJy beam $^{-1}$. The peak intensity is 174.5 mJy beam $^{-1}$ and the beam size is $1.31'' \times 1.15''$ at position angle, -87.31° . Labelling follows the conventions of Greenfield et al. (1985) and Avruch et al. (1997). The lensing galaxy is associated with radio source, G.

Fig. 3.— Spectra through the continuum-subtracted channel maps at positions corresponding to the peak of components A, B, C, D, and E (see Fig. 2). The rms map noise per channel is 0.58 ± 0.01 mJy beam $^{-1}$ and the velocity is with respect to 974 km s $^{-1}$. The last panel shows the integrated flux (mJy) measured over a region within which the continuum emission is greater than 10σ .


Fig. 4.— Map of upper limits on $f_b f_v$ (if the gas is optically thick) or on $\tau f_b f_v$ (if the gas is optically thin) for a single channel, assuming that the upper limit on ΔT_B is given by $1 \times$ the rms map noise. The map values range from a minimum of 0.00332 to a maximum of 0.385 and the greyscale ranges from 0.003 (black) to 0.06 (white). External black areas have been blanked.


Fig. 5.— Contour plot of the cloud velocity dispersion v_c (in units of km s $^{-1}$) as a function of model parameters R_c/R_* and D . Contours are as labelled. The 2.59 and 163 contours correspond to $v_c = \Delta v$ and v_c equal to the velocity range of the observations respectively.


Fig. 6.— Contour plot of $\mathcal{N}_{\min}$. Contours are at equally spaced intervals in $\log_{10}(\mathcal{N}_{\min})$. The horizontal line corresponds to the beam radius.


Fig. 7.— Contour plot of $\langle \mathcal{N}_{\text{beam}} \rangle$. Contours are at equally spaced intervals in $\log_{10}(\langle \mathcal{N}_{\text{beam}} \rangle)$. We have assumed $F_H = 1$.


Fig. 8.— Contour plot of the probability $\mathcal{P}$ for detecting a cloud as a function of the model parameters R_c/R_* and D . For this figure, we assume $F_H = 1$. Probability contours are as labelled.




















TABLE 1. Observing and Map Parameters

Observing Date	03 Jan 1997
On-source Observing time	9 ^h 54 ^m
v_{center} (ν_{center})	974 km s ⁻¹ (1415.8563 MHz)
Δv ($\Delta\nu$)	2.593 km s ⁻¹ (12.207 kHz)
No. of spectral channels	63
Bandwidth	163.4 km s ⁻¹ (769.0 kHz)
Synthesized Beam Size	1.31'' $\times$ 1.15'' @ -87.3°
Rms Map Noise/channel [Total emission]	0.61 $\pm$ 0.01 mJy beam ⁻¹
Rms Map Noise/channel [Continuum-Subtracted]	0.58 $\pm$ 0.01 mJy beam ⁻¹
Rms Map Noise [Continuum map]	0.13 mJy beam ⁻¹
T/I	404 K/(mJy beam ⁻¹)

Notes to Table 1.

All velocities heliocentric, optical definition. v_{center} is the center of the observing band. Δv gives the channel separation = the velocity resolution.